


# A roadmap for geo-neutrinos: theory and experiment




PHYSICS REPORTS  
A Review Journal of Research in Physics

**arXiv:0707.3203**

Available online at  
ScienceDirect  
www.sciencedirect.com

# Summary

- **Geo-neutrinos: a new probe of Earth's interior**
- **Open questions about radioactivity in the Earth**
- **The impact of KamLAND**
- **The potential of future experiments**
- **A possible shortcut in the roadmap**
- **(Optional?) excursions**



# Geo-neutrinos: anti-neutrinos from the Earth


U, Th and  $^{40}\text{K}$  in the Earth release heat together with anti-neutrinos, in a **well fixed ratio**:

Decay	$T_{1/2}$ [ $10^9$ yr]	$E_{\text{max}}$ [MeV]	$Q$ [MeV]	$\varepsilon_{\bar{\nu}}$ [ $\text{kg}^{-1}\text{s}^{-1}$ ]	$\varepsilon_H$ [W/kg]
$^{238}\text{U} \rightarrow ^{206}\text{Pb} + 8\ ^4\text{He} + 6e + 6\bar{\nu}$	4.47	3.26	51.7	$7.46 \times 10^7$	$0.95 \times 10^{-4}$
$^{232}\text{Th} \rightarrow ^{208}\text{Pb} + 6\ ^4\text{He} + 4e + 4\bar{\nu}$	14.0	2.25	42.7	$1.62 \times 10^7$	$0.27 \times 10^{-4}$
$^{40}\text{K} \rightarrow ^{40}\text{Ca} + e + \bar{\nu}$ (89%)	1.28	1.311	1.311	$2.32 \times 10^8$	$0.22 \times 10^{-4}$

- Earth emits (mainly) antineutrinos  $\Phi_{\bar{\nu}} \sim 10^6 \text{ cm}^{-2}\text{s}^{-1}$  whereas Sun shines in neutrinos.
- A fraction of geo-neutrinos from U and Th (not from  $^{40}\text{K}$ ) are above threshold for inverse  $\beta$  on protons:  $\bar{\nu} + \text{p} \rightarrow \text{e}^+ + \text{n} - 1.8 \text{ MeV}$
- Different components can be distinguished due to different energy spectra: e. g. anti- $\nu$  with highest energy are from Uranium.

# Probes of the Earth's interior

- Deepest hole is about 12 km
- Samples from the crust (and the upper portion of mantle) are available for geochemical analysis.
- Seismology reconstructs density profile (not composition) throughout all Earth.



## Geo-neutrinos: a new probe of Earth's interior

- ✓ They escape freely and instantaneously from Earth's interior.
- ✓ They bring to Earth's surface information about the chemical composition of the **whole** planet.




# Open questions about natural radioactivity in the Earth

**1** - What is the radiogenic contribution to terrestrial heat production?

**2** - How much U and Th in the crust?

**3** - How much U and Th in the mantle?



**4** - What is hidden in the Earth's core?  
(geo-reactor,  $^{40}\text{K}$ , ...)

**5** - Is the standard geochemical model (BSE) consistent with geo-neutrino data?



# “Energetics of the Earth and the missing heat source mystery” \*

■ Heat flow from the Earth is the equivalent of some 10000 nuclear power plants


$$H_{\text{Earth}} = (30 - 44) \text{ TW}$$

■ The BSE canonical model, based on **cosmochemical** arguments, predicts a radiogenic heat production ~ 19 TW:  
~ 9 TW **estimated** from radioactivity in the (continental) crust  
~ 10 TW **supposed** from radioactivity in the mantle  
~ 0 TW **assumed** from the core

■ Unorthodox or even heretical models have been advanced...




# Geo-ν: predictions of the BSE reference model





Signal from U+Th [TNU]	Mantovani et al. (2004)	Fogli et al. (2005)	Enomoto et al. (2005)
Pyhasalmi	51.5	49.9	52.4
Homestake	51.3		
Baksan	50.8	50.7	55.0
Sudbury	50.8	47.9	50.4
Gran Sasso	40.7	40.5	43.1
Kamioka	34.5	31.6	36.5
Curacao	32.5		
Hawaii	12.5	13.4	13.4


- **1 TNU** = one event per  $10^{32}$  free protons per year
- All calculations in agreement to the 10% level
- Different locations exhibit different contributions of radioactivity from crust and from mantle

# Geo-neutrino signal and radiogenic heat from the Earth

 region allowed by BSE: signal between 31 and 43 TNU


 region containing all models consistent with geochemical and geophysical data

 U and Th measured in the crust implies a signal at least of 24 TNU

 Earth energetics implies the signal does not exceed 62 TNU

*Fiorentini et al. (2005)*

Signal  $H(U+Th)$  from Uranium and Thorium geo-neutrinos



The graph is site dependent:

- ✓ the “slope” is universal
- ✓ the intercept depends on the site (crust effect)
- ✓ the width depends on the site (crust effect)



# KamLAND 2007 results on geo-neutrino

- In four years data ~ 630 counts in the geo- $\nu$  energy range:

- ~ 340 reactors antineutrinos

- ~ 160 fake geo- $\nu$ , from  $^{13}\text{C}(\alpha,n)$


- ~ 60 random coincidences

- ~ 70 Geo-neutrino events are obtained from subtraction.

- Adding the “Chondritic hypothesis” for U/Th:

$$N(\text{U+Th}) = 75 \pm 27$$

- This pioneering experiment has shown that the technique for identifying geo-neutrinos is now available!!!



# KamLAND 2005 results


## vs 2007

- In 2 (4) years data 152 (630) counts in the geo-neutrino energy range:

- ~ 82 (340) reactors antineutrinos

- ~ 42 (160) fake geo-neutrinos,

- ~ 5 (60) random coincidences



- Geo-neutrino events are obtained from subtraction:


$$N(\text{U+Th})=25+18 \quad \text{vs} \quad N(\text{U+Th})=75\pm 27$$

- Central value is (was 2 x) BSE prediction

\*BSE prediction is  $H(\text{U+Th}) = 16 \text{ TW}$

# Implications of KamLAND 2007 result

- The KamLAND signal  $39 \pm 15$  TNU is in perfect agreement with BSE prediction.
- It is consistent within  $1\sigma$  with:
  - Minimal model
  - Fully radiogenic model




- Concerning radiogenic heat, the 95% CL upper bound on geo-signal translates into\*  $H(U+Th) < 65$  TW

# Nuclear reactors: the enemy of geo-neutrinos


$$r = \frac{\text{Events}_{\text{reactors}}}{\text{Events}_{\text{geo } \nu}}$$

In the geo-neutrino energy window

	$r$
<b>Kamioka</b>	<b>6.7</b>
Sudbury	1.1
Gran Sasso	0.9
Pyhasalmi	0.5
Baksan	0.2
Homestake	0.2
Hawaii	0.1
Curacao	0.1





$$r = \frac{E_{\nu_{\text{react}}}}{E_{\nu_{\text{geo } \nu}}}$$



- Based on IAEA Database (2000)
- All reactors at full power

# Running and planned experiments



- Several experiments, either running or under construction or planned, have geo- $\nu$  among their goals.
- Figure shows the sensitivity to geo-neutrinos from **crust** and **mantle** together with **reactor** background.

# Borexino at Gran Sasso





- A 300-ton liquid scintillator underground detector, **running** since may 2007.

- Signal, mainly generated from the crust, is comparable to reactor background.

- From BSE expect 5 – 7 events/year\*

- In about two years should get  $3\sigma$  evidence of geo-neutrinos.

\* For 80% eff. and 300 tons  $C_9H_{12}$  fiducial mass








# SNO+ at Sudbury




- A 1000-ton liquid scintillator underground detector, obtained by replacing D<sub>2</sub>O in SNO.
- The SNO collaboration has planned to fill the detector with LS in 2009
- 80% of the signal comes from the continental crust.
- From BSE expect 28 – 38 events/year\*
- It should be capable of measuring U+Th content of the crust.

\* assuming 80% eff. and 1 kTon CH<sub>2</sub> fiducial mass



# Hanohano at Hawaii

- Project of a 10 kiloton movable deep-ocean LS detector
- ~ 70% of the signal comes from the mantle
- From BSE expect 60 – 100 events/year\*
- It should be capable of measuring U+Th content of the mantle



\* assuming 80% eff. and 10 kTon CH<sub>2</sub> fiducial mass






# LENA at Pyhasalmi

- Project of a **50 kiloton** underground liquid scintillator detector in Finland
- **80%** of the signal comes from the crust
- From BSE expect **800 – 1200** events/year\*
- LS is loaded with **0.1% Gd** which provides:
  - better neutron identification
  - moderate **directional** information

\* For  $2.5 \cdot 10^{33}$  free protons and assuming 80% eff.



*K. A. Hochmuth et al. - Astropart.Phys. 27 (2007) - arXiv:hep-ph/0509136 ; Teresa Marrodan @ Taup 2007*



# Move the mountain or the prophet?

- Geo- $\nu$  direction knows if it is coming from reactors, crust, mantle...
- Even a moderate directional information would be sufficient for source discrimination.
- $\vec{P}$  conservation implies the neutron **starts** moving “forwards”  
angle  $(\text{geo-}\nu, n) < 26^\circ$
- Directional information however is **degraded** during neutron slowing down and thermal collisions, but is not completely lost...

Geo- $\nu$  direction at Kamioka







# A shortcut in the roadmap?

- Reconstruction of geo- $\nu$  direction with Gd, Li and B loaded LS is being investigated by several groups. (See Shimizu\*, Domogatsky et al., Hochmuth et al., Poster @ TUAP 07)
- A 50 kTon 1.5%  ${}^6\text{Li}$  loaded LS in 5 years could discriminate crust and mantle contribution at the level of BSE prediction.

**A. Suzuki: "...direction measurement is the most urgent task in future geo-neutrino experiments"**




# What is needed for interpreting experimental data?

## → Geo-neutrino spectra


- The decay spectrum  $f(E)$  of geo- $\nu$  from U and Th decay chains is the input for the interaction spectrum  $s(E) = f(E) \sigma(E)$ .
- The x-section for  $I.\beta$  on free protons  $\sigma$  is known to 1% or better.
- The decay spectrum is obtained from **theoretical calculations**, assuming universal shape for allowed transitions and tabulated **branching ratios** of the transitions.



## The decay spectrum



## The interaction spectrum




The decay spectrum should be measured!

# Nuclear physics inputs needed for geo-neutrino studies \*

G Bellini<sup>1,2</sup>, G Fiorentini<sup>3,4</sup>, A Ianni<sup>5</sup>, M Lissia<sup>6</sup>, F Mantovani<sup>4,7,9</sup> and O Smirnov<sup>8</sup>



**Abstract.** Geo-neutrino studies are based on theoretical estimates of geo-neutrino spectra. We propose a method for a direct measurement of the energy distribution of antineutrinos from decays of long lived radioactive isotopes.

- Geo- $\nu$  are produced in  $\beta$  and  $\beta\gamma$  transitions



- With LS can measure the sum of energy deposited by e and  $\gamma$ .
- Anti- $\nu$  spectrum can thus be deduced from energy conservation

$$Q = E_\nu + T_e + E_\gamma$$




\*arXiv:0712.0298v1 [hep-ph]



# What is needed for interpreting experimental data?


## ➔ Regional geology

- A geochemical and geophysical study of the region (~ 200 km) around the detector is necessary for extracting the global information from the geo-neutrino signal.
- This study has been performed for Kamioka (Fiorentini et al., Enomoto et al.), it is in progress for Gran Sasso and is necessary for the other sites.






## Crustal 3D model of Central Italy





# The lesson of solar neutrinos

- ✓ Solar neutrinos started as an investigation of the solar interior for understanding **sun energetics**.
- ✓ A long and fruitful detour lead to the discovery of oscillations.
- ✓ Through several steps, we achieved a direct proof of the solar energy source, experimental solar neutrino spectroscopy, neutrino telescopes.



**The study of Earth's energetics with geo-neutrinos will also require several steps and hopefully provide surprises...**

ROAD  
WORK  
HEAD

**GAMOW 1953**  
geo- $\nu$  were born here

**KAMLAND 2005**  
1<sup>st</sup> evidence of geo- $\nu$

