

THE CLAS12 LARGE AREA RICH DETECTOR


E. Cisbani¹, M. Contalbrigo², P. Rossi³

¹ Istituto Superiore Sanita ² INFN Sezione Ferrara ³ Laboratori Nazionali di Frascati



Jefferson Lab and CEBAF


Jefferson Lab at Newport News, Virginia, USA, investigates nuclear physics via electron deep-inelastic-scattering. It uses the "Continuous Electron Beam Accelerator Facility" (CEBAF) providing a polarized electron beam with high current and energy resolution.




Hall-B

polarized H and D targets
large acceptance detector

Nucleon structure in 3D !!






CEBAF (>2013) 12 GeV electron beam



Basic Requirements

- up to 8 GeV/c momentum
- from 5° to 30° in polar angle
- π/K rejection factor > 100 (< 1% π contamination)
- non invasive: small and uniform material budget (minimize interference with Time-of-flight detector)
- minimize photon detector surface (minimize cost)
- operate in large background environment
- projective geometry (6 radial sectors, ~ 1 m gap)



Proximity focusing option:

Aerogel radiator




- 3 cm thickness
- 1.03 refraction index
- 40 mm transmission length

1 m long gap

Photon Detector:

- around 12 m² surface

K- π separation > 4 σ at 8 GeV with detector pixels smaller than 1 cm





Photon detector option with light concentrators: (a` la COMPASS)

MAPMT's:



- Pros: excellent packing factor (89%)
- Cons: non optimized for single p.e. detection






- Pros: optimized for single photon detection
- Cons: quite sizeable dead area (50%)

SiPM's:

- Pros: robust and light device
- Cons: MHz noise and high sensitivity to temperature

use narrow time window to suppress background

Test-bench at Bari



Under investigation: focusing mirror system

High momentum (> 6 GeV/c): direct light by proximity focusing

Medium momentum (3-6 GeV/c): reflected light by mirror system


Low momentum (< 3 GeV/c): time-of-flight

Detector area reduced down to ~ 4 m²

Reflecting outside

Challenge: concentrate light in a restricted area

Focusing mirrors ?



Reflecting inside

Challenge: reflected light largely adsorbed in aerogel

Partially reflective mirrors ?

