



Istituto Nazionale di Fisica Nucleare



European
Commission



Korea Institute of
Science and Technology Information

Frillion

Overview of applications of oriented crystals in accelerator physics

Dr. Alexei Sytov

sytov@fe.infn.it, alexei.sytov@kisti.re.kr

PAL Public Seminar, 23/04/03

Marie Skłodowska-Curie Action Global Individual Fellowships by A. Sytov in 2021-2024, Project TRILLION GA n. 101032975

Main goal: The **implementation** of both physics of **electromagnetic processes in oriented crystals** and the design of specific applications of crystalline effects into **Geant4** simulation toolkit as Extended Examples to bring them to a large scientific and industrial community and under a free Geant4 license.

Group:




- **A. Sytov** – project coordinator
- **L. Bandiera** – INFN supervisor
- **K. Cho** – KISTI supervisor
- **G. Kube** – DESY supervisor
- **I. Chaikovska** – IJCLab Orsay supervisor

Location:


- 2 years at **KISTI** (partner organization)
- 1 year at **INFN Section of Ferrara** (host organization)
- 1 month of secondment at **DESY** (partner organization)
- 1 month of secondment at **IJCLab Orsay** (partner organization)



How a crystal lattice looks like (from National Science Museum, Daejeon, Korea)



The world of the channeling effect




Channeling effect* of charged particles

Energies:
MeV - TeV

e⁻, e⁺, proton ...

Lindhard angle:

$$\theta < \theta_L = \sqrt{\frac{2U_0}{pv}}$$



Equivalent
magnetic field:
more than **100 T**

Channeling* is the effect of the penetration of charged particles through a monocrystal quasi parallel to its atomic axes or planes.

Planar/Axial field 10⁹/10¹¹ V/cm

*J. Stark, Zs. Phys. 13, 973–977 (1912)


J. A. Davies, J. Friesen, J. D. McIntyre, Can J. Chem. 38, 1526–1534 (1960)

M. T. Robinson, O. S. Oen, Appl. Phys. Lett. 2, 30–32 (1963)

J. Lindhard, Kgl. Dan. Vid. Selsk. Mat.-Fys. Medd. 34 No 4, 2821–2836 (1965)


Coherent effects in a crystal

Channeling radiation*




Coherent effects preserve
up to **few mrad** of particle
direction vs the crystal axis

Coherent bremsstrahlung**



Coherent pair production***






*M.A. Kumakhov, Phys. Lett. A 57(1), 17–18 (1976)

**B. Ferretti, Nuovo Cimento 7, 118 (1950).



***M. Ter-Mikaelian, Sov. Phys. JETP 25, 296 (1953).

*** H. Überall, Phys. Rev. 103, 1055 (1956).


Manufacturing and characterization of bent silicon crystals @INFN Ferrara



Piezo-activated holder




Crystal-based extraction




Planar channeling*:
Charge particle penetration through a monocrystal along its atomic planes

Crystal-based extraction/collimation



Crystal-based collimation and extraction have been used at hadron machines




Crystal-based extraction/collimation: applied only for hadrons, not yet for e-

Interesting for tens of electron synchrotrons



*J. Lindhard, Kgl. Dan. Vid. Selsk. Mat.-Fys. Medd. 34 No 4, 2821–2836 (1965)
E.N. Tsyganov, Fermilab TM-682 (1976)

Crystal-based extraction: possible setup at DESY-II



B->dipoles
QF/QD->focusing/
defocusing quadrupoles


6 GeV electrons

Advantages:

- Extraction of **primary** low-emittance and very **intense electron beam** in a **parasitic mode**.
- The **extraction line** including septum magnets already exists => **ideal for prove-of-principle**
- **Few GeV** electron beam, **typical for synchrotron light sources** existing in the world.


Applications:

- Nuclear and particle physics detectors and generic **detector R&D**
- Fixed-target experiments in **high-energy physics** including future **lepton colliders**
- Also: **crystal-based collimation** (synchrotron light sources, colliders)




Crystal-based extraction: simulation results


Coordinate distribution of extracted beam



Angular divergence of extracted beam




Extraction efficiency



Crystal parameters:

- Si (111)
- bending angle **1.75 mrad**
- Crystal length **0.175 mm**
- Crystal transverse thickness **1 cm**





Maximal extraction efficiency:
16.1 %

Where the crystal-based extraction of electrons can be applied?




Where the crystal-based extraction of electrons can be applied?




Applications*

Crystal-based collimation or beam extraction from an accelerator




Measurement of dipole magnetic and electric moments of exotic particles





Gamma-ray Space Telescope


Ultrashort crystalline calorimeter




Positron source for future e+/e- and muon colliders




Oriented crystals



Plasma acceleration




X and gamma-ray source for nuclear and medical physics




Channeling radiation in a bent crystal: Crystalline undulator

Classical scheme: magnetic undulator in a free electron laser
laser **soft X-rays** $\lambda_u \sim \text{cm}$



Innovative scheme: Crystalline undulator*
-> **Hard X-rays and gamma rays** $\lambda_u < \text{mm}$



Advantage:


- Intense X- and gamma-rays produced in a crystal, in a compact piece of material

Crystalline X and gamma-ray source **can be applied** in:

- **Nuclear physics**
- **Medical physics**

 EU project MSCA RISE N-LIGHT G. A. 872196
Coordinator MBN RESEARCH CENTER (Germany)

Crystal-based hybrid positron source*



Coherent effects in a crystal accelerate electromagnetic shower development

Advantages of the hybrid positron source:

- Higher positron yield
- Considerably lower peak deposited energy inside the target => higher beam intensities, longer target lifetime

Hybrid positron source can be applied at:

- FCC-ee
- ILC
- Muon collider

Simulation model can be also applied for ultrashort crystalline calorimeter



Plasma wake-field acceleration in oriented crystals/carbon nanotubes*

$$E[\text{GV/m}] = m_e \omega_p c / e \approx 100 \sqrt{n_0 [10^{18} \text{cm}^{-3}]}$$


**Acceleration
gradient:**

1-10 TeV/m

Considerably **higher electron density** in a **solid state** than in a gaseous plasma

Possible drive beam:



- X-rays
- electrons
- heavy high-Z beams



Channeling makes **crystal** almost **transparent** both to accelerated and to drive beam

Possible accelerated beam:

- e⁺/e⁻
- muons
- protons




* R. Ariniello, ..., and T. Tajima, **Snowmass**'2021 AF6: Advanced Acceleration Concepts, arXiv: 2203.07459

E336 collaboration future experiment at SLAC FACET-II

Channeling Acceleration in Crystals and Nanostructures and Studies of Solid Plasmas: New Opportunities

Robert Ariniello¹, Sebastien Corde², Xavier Davoine³, Henrik Ekerfelt⁴, Frederico Fiuza⁴, Max Gilljohann², Laurent Gremillet³, Yuliia Mankovska², Henryk Piekarczyk⁵, Pablo San Miguel Claveria², Vladimir Shiltsev⁵, Peter Taborek⁶, and Toshiki Tajima⁶




* R. Ariniello, ..., and T. Tajima, **Snowmass**'2021 AF6: Advanced Acceleration Concepts, arXiv: 2203.07459

T. Tajima, M. Cavenago, Crystal X-ray accelerator, Phys. Rev. Lett., 59(13), 1440 (1987).

Channeling simulation technique: Geant4 ChannelingFastSimModel


Main conception – simulation of classical trajectories of charged particles in a crystal
Multiple and single **scattering simulation** at every step



channeling*

Advantages:

- High calculation speed
- MPI parallelization for high performance computing



Baier-Katkov formula:

integration is made over the classical trajectory

$$\frac{dE}{d^3k} = \omega \frac{dN}{d^3k} \frac{\alpha}{4\pi^2} \iint dt_1 dt_2 \frac{[(E^2 + E'^2)(v_1 v_2 - 1) + \omega^2 / \gamma^2]}{2E'^2} e^{-ik'(x_1 - x_2)}$$

A.I. Sytov, V.V. Tikhomirov. NIM B 355 (2015) 383–386.

L. Bandiera, et al., Nucl. Instrum. Methods Phys. Res., Sect. B 355, 44 (2015)


*A. Sytov et al. arXiv: 2303.04385, Accepted for publication in JKPS

A. I. Sytov, V. V. Tikhomirov, and L. Bandiera. PRAB 22, 064601 (2019)

First simulations with Geant4 channeling model: beam deflection by a bent crystal


855 MeV
electrons

15 μm thick
bent crystal



volume reflection (VR)

Geant simulations vs experiment




channeling

Conclusions

- **Oriented crystals** can be **applied**:
- at **e-/e+/hadron synchrotrons** (crystal-based beam extraction/collimation)
- in **nuclear and medical physics** (radiation source)
- at e-/e+ colliders – **ILC, FCC-ee** and **muon collider** (positron source)
- Ultrahigh gradient (more than 1 **TeV/m**) **plasma wakefield acceleration**

- The goal of **TRILLION** is to implement **electromagnetic processes in oriented crystals** into **Geant4** which will bring to a large scientific and industrial community most of possible applications of a crystal.





Thank you for attention!

First Geant4 Baier-Katkov radiation model: radiation by 855 MeV electrons at Mainz Mikrotron MAMI*

G4BaierKatkov:

- **Physics list independent**
- Activated in the **DetectorConstruction** and used in **ChannelingFastSimModel**
- Can be used **outside channeling model** (e.g. in **SteppingAction**)
- Provides **radiation spectrum** for single-photon radiation mode
- Provides generation of **secondary photons**

Geant simulations vs experiment and CRYSTALRAD simulations






Electromagnetic shower acceleration

Axial field
 10^{11} V/cm





Approaching the Schwinger limit
starting from few GeV for e^+/e^-




Particle

Amorphous or randomly oriented crystal



Particle

Oriented crystal



The **radiation intensity** and the **pair production cross-section** drastically increase in **oriented crystals!**

Shower development in the field of axes is **accelerated**.
The radiation length is considerably reduced*.

KISTI-5 supercomputer NURION

Korea Institute of Science and Technology Information

Specification	KISTI-5 SKL
OS	CentOS 7.4
Processor	Intel Xeon Skylake (Gold 61148) 2.4GHz
Architecture	Multicore
Cores/CPU	20
CPUs/node	2
Cores/node	40
Total nodes	132
Total cores	5280



Our project MIRACLE, no. HP10BIW7VR Cineca Italian supercomputing center

MIRACLE, Cineca ISCRA Class B National Italian project

Medical physics and RAdiation in Crystals simuLation with gEant4

Main goal: to supply **Italian Geant4 community** and their international collaborators by CINECA HPC resources necessary to accomplish **MC_INFN** and **TRILLION** projects.

25/10/2021 - 25/01/2023

Marconi 100: 0.992 Mh for 1 year

Galileo 100: 2.4 Mh for 1 year

Italian organizations involved

- INFN Sezione di Catania
- INFN Sezione di Ferrara
- INFN Laboratori Nazionali del Sud
- INFN Napoli
- INFN Roma1
- Istituto Superiore di Sanità
- University of Messina
- University of Napoli

Foreign organizations involved

- ELI-Beamlines, Institute of Physics, (FZU), Czech Academy of Sciences
- Institute for Nuclear Problems, Belarusian State University
- University of Surrey

PI A. Sytov



How to implement an external code into Geant4?

Geant4 FastSim interface

FastSim model:

- Physics list **independent**
- Declared in the **DetectorConstruction** (just **few lines of code**)
- Is activated **only** in a **certain G4Region** at a **certain condition** and only for **certain particles**
- **Stops Geant processes** at the step of FastSim model and then resumes them

```
71  G4bool TestModel::IsApplicable(const G4ParticleDefinition& particleType)
72  {
73      return
74      &particleType == G4Proton::ProtonDefinition() ||
75      &particleType == G4AntiProton::AntiProtonDefinition() ||
76      &particleType == G4Electron::ElectronDefinition() ||
77      &particleType == G4Positron::PositronDefinition(); // ||
78      //&particleType == G4Gamma::GammaDefinition();
79  }
80
81  //.....ooo0000ooo.....ooo0000ooo.....ooo0000ooo.....ooo0000ooo.....
82
83  G4bool TestModel::ModelTrigger(const G4FastTrack& fastTrack)
84  {
102 }
103
104 //.....ooo0000ooo.....ooo0000ooo.....ooo0000ooo.....ooo0000ooo.....
105
106 void TestModel::DoIt(const G4FastTrack& fastTrack,
107                    G4FastStep& fastStep)
108 {
```

Insert particles for which the model is applicable

Insert the condition to enter the model

Insert what the model does

How to use the Geant4 channeling model in your example?

● Add to DetectorConstruction::Construct()

```
//crystal volume
G4Box* crystalSolid = new G4Box("Crystal",CrystalSizeX/2,CrystalSizeY/2,CrystalSizeZ/2.);
G4LogicalVolume* crystalLogic = new G4LogicalVolume(crystalSolid,Silicon,"Crystal");
    new G4PVPlacement(xRot,posCrystal,crystalLogic,"Crystal",logicWorld,false,0);
//crystal region (necessary for the FastSim model)
fRegion = new G4Region("Crystal");
fRegion->AddRootLogicalVolume(crystalLogic);
```

Volume declaration
(completely standard)

G4Region declaration

● Add to DetectorConstruction::ConstructSDandField()

```
void DetectorConstruction::ConstructSDandField()
{
    // ----- fast simulation -----
    //extract the region of the crystal from the store
    G4RegionStore* regionStore = G4RegionStore::GetInstance();
    G4Region* RegionCh = regionStore->GetRegion("Crystal");

    //create the channeling model for this region
    ChannelingFastSimModel* ChannelingModel = new ChannelingFastSimModel("ChannelingModel",RegionCh);
    //set the type of crystal planes
    G4String lattice = "(111)";
    //activate the channeling model
    ChannelingModel->Input(crystalLogic,lattice);
    //setting bending angle of the crystal planes (default is 0)
    BendingAngle = 0.905*mrاد;
    ChannelingModel->GetCrystalData()->SetBendingAngle(BendingAngle);

    //activate radiation model (do it only when you want to take into account
    //radiation production in an oriented crystal; it takes a lot of computational power)
    ChannelingModel->RadiationModelActivate();
}
```

Get crystal region

Channeling FastSim
model declaration

Logical volume


Model activation

Additional options

Radiation model
activation

First Geant4 channeling example for electrons/positrons

- Inspired by our experiments* of 855 MeV electron beam deflection by an ultrashort bent crystal at Mainz Mikrotron MAMI



Beam setup in **run.mac** using **GPS** commands; all the **geometry** in **DetectorConstruction**

Multithreading works! Checked at the supercomputer **Galileo100@CINECA** (Italy) **NURION@KISTI** (Korea)

Output both in **root** (only primary particles) and in **textfile** (all the particles) format



*A. Mazzolari et al. Phys. Rev. Lett. 112, 135503 (2014)


A. Sytov et al. Eur. Phys. J. C 77, 901 (2017)

How to use the Geant4 channeling model in your example?

● Add to main:

Register FastSimulationPhysics

```
G4FastSimulationPhysics* fastSimulationPhysics = new G4FastSimulationPhysics();
fastSimulationPhysics->BeVerbose();
// -- activation of fast simulation for particles having fast simulation models
// -- attached in the mass geometry:
fastSimulationPhysics->ActivateFastSimulation("e-");
fastSimulationPhysics->ActivateFastSimulation("e+");
// -- Attach the fast simulation physics constructor to the physics list:
physicsList->RegisterPhysics( fastSimulationPhysics );
```




Physics list independent


Search of MDM&EDM of short living particles using the effect of spin rotation in oriented crystals*

What we want:

- To measure **MDM** and **EDM** of exotic baryons



Experimental proof at Tevatron for Σ^{+**}



Crystal thickness must be comparable with the life distance of the particle

Possible particles:
 $\Lambda_c^+, \Xi_c^+, \dots, \tau$

* V. G. Baryshevskii, Pis'ma Zh. Tekh. Fiz. 5, 182 (1979)

**D. Chen et al. (E761 Collaboration) Phys. Rev. Lett. 69, 23 (1992)